Function Arithmetic

Just as we are able to add (+), subtract (-), multiply (·), and divide (\div) two or more numbers, we are able to +, -, ·, and \div two or more functions. Let's look at some basic notation before we go any further.

f(x) + g(x)

This means that we will add the contents of f(x) to the contents of g(x). Another way it can be written is (f + g)(x), which is more common to see. The remaining operations follow similar notation:

$$f(x) - g(x) = (f - g)(x)$$
$$f(x) \cdot g(x) = (f \cdot g)(x)$$
$$\frac{f(x)}{g(x)} = \left(\frac{f}{g}\right)(x)$$

Let's use the following two functions to demonstrate how the operations work:

$$f(x) = 3x + 2$$
 $g(x) = -2x + 1$

There are two ways to calculate values when we $+, -, \cdot$, and \div functions. (1) We can substitute a given number for x in each function first, and then perform the desired operation with the results of each function, or (2) we can perform the operation on the functions first, and then substitute the given number for x. Let's work some examples of each operation. We'll first use way (1) to find our answer in the left column, and then we will use way (2) to find our answer in the right column.

$$(f+g)(-1)$$

f(-1) = 3(-1) + 2 = -3 + 2 = -1	f(x) + g(x) = (3x + 2) + (-2x + 1)
g(-1) = -2(-1) + 1 = 2 + 1 = 3	= 3x + 2 - 2x + 1 = x + 3
f(-1) + g(-1) = -1 + 3 = 2	(f + g)(-1) = (-1) + 3 = 2

$$(g-f)(2)$$

f(2) = 3(2) + 2 = 6 + 2 = 8	g(x) - f(x) = (-2x + 1) - (3x + 2)
g(2) = -2(2) + 1 = -4 + 1 = -3	= -2x + 1 - 3x - 2 = -5x - 1
g(2) - f(2) = -3 - 8 = -11	(g-f)(2) = -5(2) - 1 = -11

$(f \cdot g)(3)$	
f(3) = 3(3) + 2 = 9 + 2 = 11	$f(x) \cdot g(x) = (3x+2)(-2x+1)$
g(3) = -2(3) + 1 = -6 + 1 = -5	$= -6x^2 + 3x - 4x + 2 = -6x^2 - x + 2$
$f(3) \cdot g(3) = 11 \cdot -5 = -55$	$(f \cdot g)(3) = -6(3)^2 - 3 + 2 = -55$

$$\left(\frac{f}{a}\right)(-3)$$

$$f(-3) = 3(-3) + 2 = -9 + 2 = -7$$

$$g(-3) = -2(-3) + 1 = 6 + 1 = 7$$

$$\frac{f(-3)}{g(-3)} = \frac{-7}{7} = -1$$

$$\left(\frac{f}{g}\right)(-3) = \frac{3(-3) + 2}{-2(-3) + 1} = \frac{-7}{7} = -1$$

Not only are we able to combine function equations together with arithmetic operations, we are able to combine function graphs together with the same operations. Consider the following two functions:



Let's find (f + g)(x). The first thing we need to do is to create a table with the ordered pairs of the points on each graph.



The next thing we do is find **matching** *x* values. For each matching *x* value we have, we perform the desired operation on the corresponding *y* values. For (f + g)(x):

